

$\Delta N_{21} = N_2 - N_1 = (0.4 \text{ to } 1.1) \times 10^{-3}$ and $\Delta N_{31} = N_3 - N_1 \approx 0.2$. Values of $\Delta N_{31} \gg \Delta N_{21}$ are common for biaxial optical crystals in both solid-state (19) and soft-matter (7, 20) systems, where typical biaxial birefringence is 10^{-3} . Experimental microscopic images, produced by optical

interference of diverging light rays traveling through a biaxial NLC (19), match the ones simulated with the use of separately measured ΔN_{21} and ΔN_{31} (Fig. 2, D to G). For n_{ω} and \tilde{S}

harvard 01161-

reveals their biaxial orientational distributions at different T and ϕ_c (Fig. 4, B to E), as well as facile electric and magnetic switching of \mathbf{n}_c (fig. S5). In the vicinity of ϕ_{cUB} , we detected only small changes of $S_c \approx -0.49$ but find that Δ_c jumps from zero to ≈ 0.6 at $\phi_c \approx 0.12\%$ (in the beginning of the two-phase coexistence) and then increases with ϕ_c to ≈ 0.95 at $\phi_c > 0.16\%$ in the biaxial phase (Fig. 4C). Orientational distribution functions are biaxial for both molecular and colloidal rods, though biaxiality of colloidal ordering is much stronger than the weak induced biaxiality in molecular ordering (Fig. 4). The biaxial NLC is not merely a superposition of two uniaxial nematic molecular and colloidal states with orthogonal ordering directions. Anisotropic molecular interactions at interfaces of orientationally ordered nanorods lift the uniaxial symmetry of molecular order and, simultaneously, induce biaxiality in the distribution of colloidal nanorod orientations. Tensorial order parameters describing this behavior of molecules and colloids, $\mathbf{Q}_{m,c} = \text{diag}\{S_{m,c}, (\Delta_{m,c} - S_{m,c})/2, -(\Delta_{m,c} + S_{m,c})/2\}$, determine physical properties—such as the optical anisotropies characterized in Figs. 1 and 2—



Hybrid molecular-colloidal liquid crystals

Haridas Mundoor, Sungoh Park, Bohdan Senyuk, Henricus H. Wensink and Ivan I. Smalyukh

Science **360** (6390), 768-771.
DOI: 10.1126/science.aap9359

Finding order in twos

In nematic liquid crystals, the local orientation of the molecules hovers around an average direction. The orientational control bestows unusual optical properties. In theory, with the right sort of two-dimensional shape, it should be possible to create nematics with biaxial ordering, but this has proven elusive. Mundoor *et al.* dispersed colloidal rods into a nematic solvent (see the Perspective by Poulin). Within a range of temperature and concentration, the rods ordered orthogonally to the solvent molecules, thus giving the mixture the type of properties that one would expect from a biaxial liquid crystal.

Science, this issue p. 768; see also p. 712

ARTICLE TOOLS	http://science.sciencemag.org/content/360/6390/768
SUPPLEMENTARY MATERIALS	http://science.sciencemag.org/content/suppl/2018/05/16/360.6390.768.DC1
RELATED CONTENT	http://science.sciencemag.org/content/sci/360/6390/712.full
REFERENCES	This article cites 27 articles, 3 of which you can access for free http://science.sciencemag.org/content/360/6390/768#BIBL
PERMISSIONS	http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2018 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works